


Syphilis


Sexually Transmitted Disease Surveillance 1998

Division of STD Prevention


Syphilis - Reported cases by stage of illness: United States, 1941-1998



Primary and secondary syphilis - Reported rates: United States, 1970-1998 and the Healthy People year 2000 objective




Primary and secondary syphilis - Rates by state: United States and outlying areas, 1998




Note: The total rate of primary and secondary syphilis for the United States and outlying areas (including Guam, Puerto Rico and Virgin Islands) was 2.6 per 100,000 population. The Healthy People year 2000 objective is 4.0 per 100,000 population.

Primary and secondary syphilis - Counties with rates above and
counties with rates below the Healthy People year 2000 objective:
United States, 1998




Primary and secondary syphilis - Rates by region: United States, 1981-1998 and the Healthy People year 2000 objective




Primary and secondary syphilis - Rates by urban-rural category and geographic region, 1998


Rate (per 100,000 population)




Primary and secondary syphilis - Rates in selected U.S. Cities of >200,000 population, 1981-1998 and the Healthy People year 2000 objective



Primary and secondary syphilis - Rates by gender: United States, 1981-1998 and the Healthy People year 2000 objective




Primary and secondary syphilis - Rates by race and ethnicity: United States, 1981-1998 and the Healthy People year 2000 objective




Note: "Other" includes Asian/Pacific Islander and American Indian/Alaska Native populations. Black, White, and Other are non-Hispanic.

Primary and secondary syphilis - Age- and gender-specific rates: United States, 1998




Congenital syphilis - Reported cases for infants <1 year of age and rates of primary and secondary syphilis among women: United States, 1970-1998



Note: The surveillance case definition for congenital syphilis changed in 1988

Congenital syphilis - Rates for infants <1 year of age: United States, 1981-1998 and the Healthy People year 2000 objective



Note: The surveillance case definition for congenital syphilis changed in 1988